



**STAND UP
TO PAH**

TAKE BACK MORE GOOD DAYS with Orenitram®

Proven to help slow the progression of PAH, improve exercise capacity, and may help reduce PAH symptoms.

IMPORTANT SAFETY INFORMATION FOR ORENITRAM

Who should not take Orenitram?

Do not take Orenitram if you have severe liver problems.

What should I tell my healthcare provider before taking Orenitram?

Tell your healthcare provider:

- If you have liver problems or diverticulosis.

Please see **Full Prescribing Information** and **Patient Information** at www.orenitram.com.

PAH=pulmonary arterial hypertension.



orenitram®
treprostiniil

EXTENDED-RELEASE TABLETS

PAH and the body

PAH symptoms happen when the blood vessels inside your lungs become narrow, making it harder for blood to flow. This can be caused by imbalances of certain natural substances in your body: **endothelin, nitric oxide, activin, and prostacyclin.**



Oral prostacyclin provides another approach to treating PAH

Orenitram helps to replace the natural prostacyclin you may be missing. It's formulated to help keep the blood vessels in the lungs open, which makes it easier for your heart to pump blood through your lungs.



Is your current PAH therapy falling short of your treatment goals?

Orenitram targets a different pathway from other oral medicines you may be taking for PAH. That's why adding it to your treatment plan may be the next step to help you do more to control your PAH.

IMPORTANT SAFETY INFORMATION FOR ORENITRAM (CONT'D)

What should I tell my healthcare provider before taking Orenitram?

Tell your healthcare provider:

- If you are pregnant, breastfeeding, and/or plan to become pregnant or breastfeed. It is not known if Orenitram will harm your unborn baby or if Orenitram passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with Orenitram.

Meet Orenitram

Orenitram is a prescription medicine used to treat PAH. It helps slow down the progression of your disease and improve your ability to exercise.

20+
YEARS

Did you know?

Orenitram is the **tablet form of the same proven type of medicine** that has been available as a pump therapy for more than 20 years!



Not shown at actual size.

IMPORTANT SAFETY INFORMATION FOR ORENITRAM (CONT'D)

What should I tell my healthcare provider before taking Orenitram?

Tell your healthcare provider:

- **About all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Orenitram and other medicines may affect each other causing side effects. Do not start any new medicine until you check with your healthcare provider. Especially tell your healthcare provider if you take another medicine that contains treprostinil, such as Remodulin® or Tyvaso®.

Please see [Full Prescribing Information](#) and [Patient Information](#) at www.orenitram.com.


orenitram[®]
treprostinil
EXTENDED-RELEASE TABLETS

Why Orenitram?

Orenitram may help you reach your treatment goals and control your PAH. It has been proven to delay the progression of PAH and may help improve PAH symptoms. **Many people feel results in as few as 3 months.***

Orenitram may help you:



Increase your **ability to walk further**[†]



Decrease **shortness of breath** during daily activities*



Reduce the **strain** on your heart*

*In a clinical study that included 690 patients who were all taking a single oral PAH medicine (PDE-5i, sGCS, or ERA). At the beginning of the study, these patients added either Orenitram or placebo.

†In a clinical study that included 349 patients not taking any PAH therapy. At the beginning of the study, these patients began taking either Orenitram or placebo.

IMPORTANT SAFETY INFORMATION FOR ORENITRAM (CONT'D)

How should I take Orenitram?

- **Do not change your dose or suddenly stop taking Orenitram without first talking to your healthcare provider.**
- Orenitram is usually taken 3 times a day (about every 8 hours) or 2 times a day (about every 12 hours). Your healthcare provider will tell you how often you should take Orenitram. If you have side effects, your healthcare provider may tell you to change your dose or when you take Orenitram. Take Orenitram with food.

Please see [Full Prescribing Information](#) and [Patient Information](#) at www.orenitram.com.



9 out of 10

PATIENTS IMPROVED OR MAINTAINED

their risk status with Orenitram—having a lower risk status may help improve outcomes in some patients*

14.7% of patients taking Orenitram improved and 74% maintained WHO FC at week 24 vs 10.8% improved and 70.9% maintained in the placebo group.

*In a clinical study that included 690 patients who were all taking a single oral PAH medicine (PDE-5i, sGCS, or ERA). At the beginning of the study, these patients added either Orenitram or placebo.



IMPORTANT SAFETY INFORMATION FOR ORENITRAM (CONT'D)

How should I take Orenitram?

- Swallow Orenitram tablets whole. Do not split, chew, crush, or break your Orenitram tablets. Do not take Orenitram tablets that are damaged or broken. **If Orenitram tablets are not taken whole, they may release too much medicine at one time. This can lead to side effects.**
- If you miss your dose of Orenitram, take the missed dose as soon as possible with food.
- If you miss 2 or more doses of Orenitram, call your healthcare provider to see if you need to change your dose.



orenitram[®]
treprostinil
EXTENDED-RELEASE TABLETS

What is titration?

When you start Orenitram, your healthcare team will work with you to gradually increase your dose—a **process called titration**.

During this time, your body is still getting used to the medicine, and you will likely experience side effects. Remember that side effects may get better with time as you continue increasing your dose of Orenitram.

Why is titration important?

Titration is all about finding the most effective dose for you.

That's why it's important to stay committed to your treatment. You may not get the most benefit from Orenitram until you reach your target dose.



IMPORTANT SAFETY INFORMATION FOR ORENITRAM (CONT'D)

How should I take Orenitram?


- If you take too much Orenitram, call your healthcare provider or go to the nearest hospital emergency room right away.
- You may see the tablet shell in your stools (bowel movements). This is usually normal. The tablet shell is not digested. If you have diverticulosis, the tablet shell may get stuck in a blind pouch or diverticulum in your intestine.

Personalized dosing

Since PAH is a progressive disease, your treatment needs may change over time. Your healthcare provider can continue to increase or change your dose to meet your needs. That way, you may be able to stay with Orenitram for the long term to help control your PAH.*

5 DIFFERENT TABLET STRENGTHS

make it easier for your healthcare provider to adjust your dose.



*The recommended maximum daily dose is 120 mg.

IMPORTANT SAFETY INFORMATION FOR ORENITRAM (CONT'D)

What are the possible side effects of Orenitram?

Orenitram can cause serious side effects, including worsening of PAH symptoms.

- Stopping Orenitram suddenly may cause worsening of your PAH symptoms. **Do not change your dose or suddenly stop taking Orenitram without first talking to your healthcare provider.**

Please see [Full Prescribing Information](#) and [Patient Information](#) at www.orenitram.com.


orenitram[®]
treprostinil
EXTENDED-RELEASE TABLETS

Taking Orenitram

Like other medicines you may be taking, Orenitram is an oral tablet. **You take it with food, usually 3 times daily (every 8 hours).*** Talk to your healthcare team about food choices that may work best for you.



6 am breakfast • 2 pm afternoon snack • 10 pm bedtime snack

Need some snack suggestions for taking your Orenitram dose?

You can find yummy recipes and snack ideas, as well as an interactive food guide, on orenitram.com.

*Orenitram may also be taken 2 times a day (every 12 hours). Tablet strength may be increased depending on how well you tolerate treatment.



IMPORTANT SAFETY INFORMATION FOR ORENITRAM (CONT'D)

What are the possible side effects of Orenitram?

Orenitram can cause serious side effects, including worsening of PAH symptoms.

- **The most common side effects of Orenitram include** headache, diarrhea, nausea, vomiting, flushing, and pain in arms, legs, and jaw. These are not all of the possible side effects of Orenitram. Tell your healthcare provider if you have any side effect that bothers you or does not go away.
- Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at www.fda.gov/MedWatch or call [1-800-FDA-1088](tel:1-800-FDA-1088).

Tips for staying on track



Taking Orenitram with food helps the body absorb the medicine and may help reduce side effects.

But you don't have to take it with a full meal—a snack is fine




To avoid missing a dose, try adding alerts to your calendar or smart phone alarm



If you've started on Orenitram and you miss a dose, take the dose as soon as possible with food. If you miss 2 or more doses, call your healthcare provider for guidance

Hear about Jenny's transition to Orenitram and the positive changes to her lifestyle. [Click here](#) to watch video.



IMPORTANT SAFETY INFORMATION FOR ORENITRAM (CONT'D)

How should I take Orenitram?

- You may see the tablet shell in your stools (bowel movements). This is usually normal. The tablet shell is not digested. If you have diverticulosis, the tablet shell may get stuck in a blind pouch or diverticulum in your intestine.

Please see [Full Prescribing Information](#) and [Patient Information](#) at www.orenitram.com.


orenitram[®]
treprostiniil
EXTENDED-RELEASE TABLETS

Expected side effects

The most common side effects of Orenitram include headache, diarrhea, nausea, vomiting, flushing, and pain in the arms, legs, and jaw.

- Side effects are a normal part of taking Orenitram, especially at first
- They may get better with time as you continue increasing your dose of Orenitram
- Taking Orenitram at the same time you eat or 15 minutes after may help minimize side effects

It pays to be proactive

Side effects are a reality—but that's not the whole story. Write down your experience. Talk to your doctor. And know that feeling better is possible if you stick with it.



IMPORTANT SAFETY INFORMATION FOR ORENITRAM (CONT'D)

What are the possible side effects of Orenitram?

Orenitram can cause serious side effects, including worsening of PAH symptoms.

- Stopping Orenitram suddenly may cause worsening of your PAH symptoms. **Do not change your dose or suddenly stop taking Orenitram without first talking to your healthcare provider.**

Please see [Full Prescribing Information](#) and [Patient Information](#) at www.orenitram.com.

Managing side effects

The more you communicate with your healthcare team about side effects, the better. That way, they can suggest some ways to help:



Headaches

Over-the-counter treatments such as acetaminophen or ibuprofen



Nausea & vomiting

Various anti-nausea and anti-vomiting medications



Diarrhea

Adding fiber to your diet or taking anti-diarrhea medication

United Therapeutics does not provide medical advice. Your healthcare provider will help you to develop a plan to manage side effects.

IMPORTANT SAFETY INFORMATION FOR ORENITRAM (CONT'D)

What are the possible side effects of Orenitram?

Orenitram can cause serious side effects, including worsening of PAH symptoms.

- **The most common side effects of Orenitram include** headache, diarrhea, nausea, vomiting, flushing, and pain in arms, legs, and jaw. These are not all of the possible side effects of Orenitram. Tell your healthcare provider if you have any side effect that bothers you or does not go away.



Staying focused on your treatment goals

Because of the way Orenitram works, you'll probably experience side effects before any benefits. It can be tough.

But staying on track means you have a chance to see real improvements in your PAH symptoms. Try these things to help keep your goal in focus:

10
years

Remember that you are not alone

For over 10 years, Orenitram has helped many patients do more to control their PAH.



Lean on your community

Be open with friends and family about what you're going through—and don't be afraid to reach out for support. Online communities can be great too.



Know what you want

What is PAH getting in the way of? Write a list of hopes and goals for life with Orenitram.



Keep in touch with your healthcare team

It is important to talk to your healthcare provider about side effects you experience so you can get advice on how to help manage them.

IMPORTANT SAFETY INFORMATION FOR ORENITRAM (CONT'D)

What are the possible side effects of Orenitram?

Orenitram can cause serious side effects, including worsening of PAH symptoms.

- Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at www.fda.gov/MedWatch or call 1-800-FDA-1088.

Support every step of the way

If you and your healthcare provider decide that Orenitram may help you reach your treatment goals, United Therapeutics, the maker of Orenitram, provides support at every step in the process.

United Therapeutics Cares: United for you

United Therapeutics Cares provides you with ongoing, one-on-one education and support beginning on day one of your United Therapeutics treatment journey. Your dedicated Patient Navigator will personalize support and answer questions about what to expect as you begin treatment, insurance coverage, out-of-pocket cost assistance and prescription fulfillment.



Work with your provider to enroll in United Therapeutics Cares. Visit [UnitedTherapeuticsCares.com](https://www.unitedtherapeuticscares.com) to learn more.

If you have any questions, call [1-844-864-8437](tel:1-844-864-8437) to speak with your Patient Navigator, Monday through Friday, 8:30 am – 7 pm ET.

Get to know your Specialty Pharmacy

Your Specialty Pharmacy is an essential partner in your treatment journey. They can help with access to medication and supplies, in-home training, follow-up visits, and a 24-hour hotline.

United Therapeutics works with 2 Specialty Pharmacies:

Accredo
By EVERNORTH

www.ACCREDO.com
1-866-344-4874

CVS specialty™

www.CVSSPECIALTY.com
1-877-242-2738



orenitram®
treprostiniL
EXTENDED-RELEASE TABLETS

Please see [Full Prescribing Information](#) and [Patient Information](#) at www.orenitram.com.

Get to know MyOrenitram.com

For an experience thoughtfully created to meet your needs, visit MyOrenitram.com. There, you'll find support for your specific dosing schedule, tips to help you along the way, and you can listen to stories from patients like you.

[Watch this video](#) to learn more about [MyOrenitram.com](#)



Learn about possible cost savings

With our co-pay program, eligible patients may pay as little as \$0 per prescription for Orenitram.*

Learn more at www.UTcopay.com



*TO ENROLL IN THIS PROGRAM, YOU MUST UNDERSTAND AND AGREE TO COMPLY WITH THE ELIGIBILITY REQUIREMENTS AND TERMS OF USE.

Eligibility requirements for this program are: Patients must be 18 years or older to use this Program. The Program is valid only for patients with commercial (also known as private) insurance who are taking the medication for an FDA-approved indication. Patients using Medicare, Medicaid, or any other state or federal government program to pay for their medications are not eligible. Patients who start utilizing government coverage during the term of the Program will no longer be eligible. Eligible patients must be residents of the United States or Puerto Rico.

For full program details and Terms and Conditions, visit www.UTcopay.com.

Please see [Full Prescribing Information](#) and [Patient Information](#) at www.orenitram.com.

Join Orenitram Connect

Sign up to learn more about adding Orenitram to your PAH treatment plan. With Orenitram Connect, you'll also receive lifestyle tips, helpful emails, and other useful tools.



Connect with us at
www.orenitram.com/sign-up

Get social

The Orenitram Instagram, Facebook, and YouTube pages provide helpful resources, videos, and support from others, like you.



IMPORTANT SAFETY INFORMATION FOR ORENITRAM (CONT'D)

The risk information provided here is not comprehensive. To learn more about Orenitram, talk with your healthcare provider.

Please see [Full Prescribing Information](#) and [Patient Information](#) at www.orenitram.com or call Customer Service at 1-877-UNITHER (1-877-864-8437).



orenitram[®]
treprostinil
EXTENDED-RELEASE TABLETS

IMPORTANT SAFETY INFORMATION FOR ORENITRAM

What is Orenitram?

Orenitram is a prescription medicine used to treat pulmonary arterial hypertension (PAH) which is high blood pressure in the arteries of your lungs. Orenitram can help slow down the progression of your disease and improve your ability to exercise. It is not known if Orenitram is safe and effective in children.

Who should not take Orenitram?

Do not take Orenitram if you have severe liver problems.

What should I tell my healthcare provider before taking Orenitram?

Tell your healthcare provider:

- If you have liver problems or diverticulosis.
- If you are pregnant, breastfeeding, and/or plan to become pregnant or breastfeed. It is not known if Orenitram will harm your unborn baby or if Orenitram passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby during treatment with Orenitram.
- **About all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Orenitram and other medicines may affect each other causing side effects. Do not start any new medicine until you check with your healthcare provider. Especially tell your healthcare provider if you take another medicine that contains trestatinil, such as Remodulin® or Tyvaso®.

How should I take Orenitram?

- **Do not change your dose or suddenly stop taking Orenitram without first talking to your healthcare provider.**
- Orenitram is usually taken 3 times a day (about every 8 hours) or 2 times a day (about every 12 hours). Your healthcare provider will tell you how often you should take Orenitram. If you have side effects, your healthcare provider may tell you to change your dose or when you take Orenitram. Take Orenitram with food.

- Swallow Orenitram tablets whole. Do not split, chew, crush, or break your Orenitram tablets. Do not take Orenitram tablets that are damaged or broken. **If Orenitram tablets are not taken whole, they may release too much medicine at one time. This can lead to side effects.**
- If you miss your dose of Orenitram, take the missed dose as soon as possible with food.
- If you miss 2 or more doses of Orenitram, call your healthcare provider to see if you need to change your dose.
- If you take too much Orenitram, call your healthcare provider or go to the nearest hospital emergency room right away.
- You may see the tablet shell in your stools (bowel movements). This is usually normal. The tablet shell is not digested. If you have diverticulosis, the tablet shell may get stuck in a blind pouch or diverticulum in your intestine.

What are the possible side effects of Orenitram?

Orenitram can cause serious side effects, including worsening of PAH symptoms.

- Stopping Orenitram suddenly may cause worsening of your PAH symptoms. **Do not change your dose or suddenly stop taking Orenitram without first talking to your healthcare provider.**
- **The most common side effects of Orenitram include** headache, diarrhea, nausea, vomiting, flushing, and pain in arms, legs, and jaw. These are not all of the possible side effects of Orenitram. Tell your healthcare provider if you have any side effect that bothers you or does not go away.
- Call your healthcare provider for medical advice about side effects. You may report side effects to the FDA at www.fda.gov/MedWatch or call 1-800-FDA-1088.

The risk information provided here is not comprehensive. To learn more about Orenitram, talk with your healthcare provider.

Please see [Full Prescribing Information](#) and [Patient Information](#) at www.orenitram.com or call Customer Service at 1-877-UNITHER (1-877-864-8437).

OREISconOCT19

Orenitram is a registered trademark of United Therapeutics Corporation.

All other brands are trademarks of their respective owners. The makers of these brands are not affiliated with and do not endorse United Therapeutics or its products.

© 2024 United Therapeutics Corporation. All rights reserved. US/ORE/0773 Printed in USA.

